

Stat 3011 Midterm 1 (Class Part)

Problem 1

All of this is summarized in the box on p. 25 in Wild and Seber.

On the first issue (what is the difference),

- In an **experiment**, the experimenter *determines* which experimental units receive which treatments (ideally by randomized assignment to treatment groups).
- In an **observational study**, which subject gets which treatment is outside the experimenter's control. Perhaps the subjects or their doctors choose. Perhaps it just happens (exposure to a toxic substance in the environment, for example).

On the second issue (what is the implication)

- A properly designed and executed **experiment** can reliably demonstrate *causation*.
- An **observational study** can only suggest possible causes. It *cannot* reliably establish *causation*. A properly designed and executed experiment is needed to follow up results implied by an observational study.

Problem 2

(a) The mean is

$$\begin{aligned}\sum_x x \operatorname{pr}(x) &= 1 \cdot \frac{1}{7} + 2 \cdot \frac{1}{7} + 3 \cdot \frac{1}{7} + 4 \cdot \frac{1}{7} + 5 \cdot \frac{1}{7} + 6 \cdot \frac{1}{7} + 7 \cdot \frac{1}{7} \\ &= \frac{1 + 2 + 3 + 4 + 5 + 6 + 7}{7} \\ &= \frac{28}{7} \\ &= 4\end{aligned}$$

(b) The standard deviation is

$$\operatorname{sd}(X) = \sqrt{E\{(X - \mu)^2\}}$$

where μ is the mean calculated in part (a), and

$$\begin{aligned}
 E\{(X - \mu)^2\} &= \sum_x (x - \mu)^2 \text{pr}(x) \\
 &= (1 - 4)^2 \cdot \frac{1}{7} + (2 - 4)^2 \cdot \frac{1}{7} + (3 - 4)^2 \cdot \frac{1}{7} + (4 - 4)^2 \cdot \frac{1}{7} \\
 &\quad + (5 - 4)^2 \cdot \frac{1}{7} + (6 - 4)^2 \cdot \frac{1}{7} + (7 - 4)^2 \cdot \frac{1}{7} \\
 &= \frac{(-3)^2 + (-2)^2 + (-1)^2 + 0^2 + 1^2 + 2^2 + 3^2}{7} \\
 &= \frac{9 + 4 + 1 + 0 + 1 + 4 + 9}{7} \\
 &= \frac{28}{7} \\
 &= 4
 \end{aligned}$$

Thus $\text{sd}(X) = \sqrt{4} = 2$.

Problem 3

(a) By the multiplication rule (using the assumed statistical independence)

$$\text{pr}(\text{no accidents in 30 days}) = \text{pr}(\text{no accident in one day})^{30}$$

So in order to answer this question we have to first answer the subsidiary question: what is the latter probability? By the complement rule

$$\begin{aligned}
 \text{pr}(\text{no accident in one day}) &= 1 - \text{pr}(\text{an accident in one day}) \\
 &= 1 - 0.002 \\
 &= 0.998
 \end{aligned}$$

Thus

$$\text{pr}(\text{no accidents in 30 days}) = 0.998^{30} = 0.941708$$

(b) The events in parts (a) and (b) of this problem are complementary, so by the complement rule each is one minus the other

$$\text{pr}(\text{at least one accident in 30 days}) = 1 - 0.941708 = 0.058292$$

Problem 4

Curve A: skewed, long right tail, unimodal.

Curve B: symmetric, unimodal.

Curve C: skewed, long right tail, bimodal.

Curve D: symmetric, bimodal.