Stat 5102 (Geyer) Spring 2012 Homework Assignment 3 Due Wednesday, February 8, 2012

Solve each problem. Explain your reasoning. No credit for answers with no explanation. If the problem is a proof, then you need words as well as formulas. Explain why your formulas follow one from another.

3-1. Show that the family of $Gam(\alpha, \lambda)$ distributions with α known and λ unknown, so the parameter space is

$$\{\lambda \in \mathbb{R} : \lambda > 0\}$$

is a scale family.

- **3-2.** Suppose S_n^2 is the sample variance calculated from an IID normal random sample of size n.
- (a) Calculate the bias of S_n as an estimator of the population standard deviation σ .
- (b) Find the constant a such that aS_n has the smallest mean square error as an estimator of σ .
- **3-3.** Suppose U and V are statistics that are independent random variables and both are unbiased estimators of a parameter θ . Write $\text{var}(U) = \sigma_U^2$ and $\text{var}(V) = \sigma_V^2$, and define another statistic T = aU + (1-a)V where a is an arbitrary but known constant.
- (a) Show that T is an unbiased estimator of θ .
- (b) Find the a that gives T the smallest mean square error.
- **3-4.** The slides don't give any examples of estimators that are *not* consistent. Give an example of an inconsistent estimator of the population mean.
- **3-5.** If $X \sim \text{Bin}(n, p)$, show that $\hat{p}_n = X/n$ is a consistent and asymptotically normal estimator of p, and give the asymptotic distribution of \hat{p}_n .
- **3-6.** If X_1, X_2, \ldots are IID from a distribution having a variance σ^2 , show that both V_n and S_n^2 are consistent estimators of σ^2 .

- **3-7.** Suppose X_1, X_2, \ldots are IID Geo(p).
- (a) Find a method of moments estimator for p.
- (b) Find the asymptotic normal distribution of your estimator.
- **3-8.** Suppose X_1, X_2, \ldots are IID Beta $(\alpha, 2)$.
- (a) Find a method of moments estimator for α .
- (b) Find the asymptotic normal distribution of your estimator.
- **3-9.** Let X_1, X_2, \ldots, X_n be an i. i. d. sample from a Beta (θ, θ) model, where θ is an unknown parameter. Find a method of moments estimator of θ .

Review Problems from Last Year's Tests

3-10. For the following data

$$1.5 \quad 2.0 \quad 2.5 \quad 3.0 \quad 4.5$$

- (a) Find the mean of the empirical distribution.
- (b) Find the median of the empirical distribution.
- (c) Find $P_n(X \leq 3)$ under the empirical distribution.
- (d) Find the 0.25 quantile of the empirical distribution.
- **3-11.** Find the asymptotic distribution of the sample median of an IID sample from the $\text{Exp}(\lambda)$ distribution.
- **3-12.** Suppose X_1, X_2, \ldots are IID NegBin(r, p), where r is known and p is unknown.
- (a) Find a method of moments estimator for p.
- (b) Find the asymptotic normal distribution of your estimator.