

Joint analysis of survival and reproduction over 10 years in perennial *Echinacea* plants from seven populations

Stuart Wagenius¹, Ruth Shaw², Charles Geyer²

¹Chicago Botanic Garden

²University of Minnesota

<http://www.stat.umn.edu/geyer/aster/>

<http://echinacea.umn.edu/>






H. Hangelbroek

Echinacea: Model prairie species

- **Asteraceae**
- **Herbaceous**
- **Long-lived**
 - juvenile
 - adult
- **Reproduces by seed**





1.6 km

Common Garden

571 plugs planted in
1996.

Randomized
positions on 1m x 1m
grid.


Old field with
competition.

Biennial spring burns.

Annual
measurements &
harvests.



Survival, 1996-2006



1



2

3

4

5

6



Problems measuring fitness

1. Each fitness component for a given individual is conditional on the individual's state for an earlier component of fitness.

1





2

3


4

5

6



Histogram of fruit counts



Problems measuring fitness

1. Each fitness component for a given individual is conditional on the individual's state for an earlier component of fitness.
2. No single probability distribution is suitable for modeling all components of fitness.



Year

1

2

3

4

5

6

YES → YES → YES → YES → YES → YES

**Survive
to year?**

Binomial

Year 1 2 3 4 5 6

**Flower?
Binomial**

NO

YES

YES

NO

YES



YES → YES → YES → YES → YES → YES

**Survive
to year?
Binomial**

Year 1 2 3 4 5 6


**Head count,
Poisson**

0 1 2 0 5



**Flower?
Binomial**

NO YES YES NO YES



YES → YES → YES → YES → YES → YES

**Survive
to year?
Binomial**

Year 1 2 3 4 5 6


**Head count,
Poisson**

0 1 2 0 0



**Flower?
Binomial**

NO YES YES NO NO



YES → YES → YES → YES → NO → NO

**Survive
to year?
Binomial**

*

problem

solution

Measuring

Echinacea project

echinacea.umn.edu

Modeling


aster life history analysis

www.stat.umn.edu/geyer/aster/

Implementing


R statistical software

www.r-project.org




Proportion of plants flowering

(& most heads per plant each year)



Common Garden 2006



Model Comparisons

Analysis of Deviance Table

Model 1: $\text{resp} \sim \text{varb} + \text{level}:(\text{nsloc} + \text{ewloc})$

Model 2: $\text{resp} \sim \text{varb} + \text{level}:(\text{nsloc} + \text{ewloc}) + \text{hdct} * \text{pop} - \text{pop}$


Model 3: $\text{resp} \sim \text{varb} + \text{level}:(\text{nsloc} + \text{ewloc}) + \text{hdct} * \text{pop}$

Model 4: $\text{resp} \sim \text{varb} + \text{level}:(\text{nsloc} + \text{ewloc}) + \text{level} * \text{pop}$

Model No.	Model d.f.	Model Deviance	Test d.f.	Test Deviance	Test p -value	.
1	31	5308.1				
2	37	5293.8	6	14.3	0.026	
3	43	5266.9	6	26.9	0.00015	
4	49	5255.7	6	11.1	0.1	

Total plant fitness 1996-2006


Total number of flower heads per plant



Year 1 2 3 4 5 6

**Rosettes,
Poisson**

 1 5 6 8 11




YES → YES → YES → YES → YES → YES

**Survive
to year?
Binomial**

Rosettes per plant, 2003-2006

unconditional mean value parameter



Conclusions

Population differences in head count (fitness).

Population differences in rosette count (biomass).

The new statistical model "aster" jointly models composite sequential responses with predictors in a single analysis.

Further study with Echinacea

Fruit counts

179490 & counting

More pops & pop characteristics

common garden & remnants

Experimental inbreeding



A big plant

Other aster applications

Selection experiments in annuals

Herbivory on inbred plants

Aphid population growth from life tables

What data do you have?

<http://www.stat.umn.edu/geyer/aster/>

Acknowledgements

Co-authors

Gretel Kiefer

Field assistants

Echinacea project

echinacea.umn.edu

Pendragon & PalmOS

NSF: DMS-0083468

& DEB-0545072

R project

www.r-project.org



